

Write your name here			
Surname MODEL ANSWERS	Other names		
Pearson Edexcel International Advanced Level	Centre Number 	Candidate Number 	
Core Mathematics C3 Advanced			
Monday 27 January 2014 – Morning Time: 1 hour 30 minutes		Paper Reference 6665A/01	
You must have: Mathematical Formulae and Statistical Tables (Pink)			Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P43136A

©2014 Pearson Education Ltd.
5/5/5/5/



PEARSON

1.

$$f(x) = \sec x + 3x - 2, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}$$

(a) Show that there is a root of $f(x) = 0$ in the interval $[0.2, 0.4]$

(2)

(b) Show that the equation $f(x) = 0$ can be written in the form

$$x = \frac{2}{3} - \frac{1}{3\cos x}$$

(1)

The solution of $f(x) = 0$ is α , where $\alpha = 0.3$ to 1 decimal place.(c) Starting with $x_0 = 0.3$, use the iterative formula

$$x_{n+1} = \frac{2}{3} - \frac{1}{3\cos x_n}$$

to calculate the values of x_1 , x_2 and x_3 , giving your answers to 4 decimal places.

(3)

(d) State the value of α correct to 3 decimal places.

(1)

$$a) f(0.2) = \sec(0.2) + 3(0.2) - 2$$

$$= \frac{1}{\cos(0.2)} + 3(0.2) - 2$$

$$= -0.37966$$

$$= \underline{-0.380} \quad (3 \text{ sig fig})$$

$$f(0.4) = \frac{1}{\cos(0.4)} + 3(0.4) - 2$$

$$= +0.28570$$

$$= \underline{+0.286} \quad (3 \text{ sig fig})$$

Sign change in the interval $[0.2, 0.4] \Rightarrow$ root,
function continuous.



Question 1 continued

$$b) f(x) = \sec x + 3x - 2 = 0$$

$$3x = 2 - \frac{1}{\cos x}$$

$$x = \frac{2}{3} - \frac{1}{3 \cos x}$$

$$c) \quad 0.3 = \text{''} \quad \text{''} \quad (\text{ANS}) = \text{''} = \text{''} = \text{''}$$

$$x_{n+1} = \frac{2}{3} - \frac{1}{3 \cos x_n}$$

$$x_0 \quad 0.3 \quad (\text{to 4 D.P.})$$

$$x_1 \quad 0.3177494662 \quad 0.3177 \text{ rad.}$$

$$x_2 \quad 0.3157676649 \quad 0.3158 \text{ rad.}$$

$$x_3 \quad 0.3159955441 \quad 0.3160 \text{ rad.}$$

$$d) \quad \underline{\alpha = 0.316 \text{ radians}} \quad (3 \text{ D.P.})$$

Q1

(Total 7 marks)



P 4 3 1 3 6 A 0 3 2 8

2.

$$f(x) = \frac{15}{3x+4} - \frac{2x}{x-1} + \frac{14}{(3x+4)(x-1)}, \quad x > 1$$

(a) Express $f(x)$ as a single fraction in its simplest form.

(4)

(b) Hence, or otherwise, find $f'(x)$, giving your answer as a single fraction in its simplest form.

(3)

a)

$$f(x) = \frac{15(x-1)}{(3x+4)(x-1)} - \frac{2x(3x+4)}{(3x+4)(x-1)} + \frac{14}{(3x+4)(x-1)}$$

$$= \frac{15x - 15 - 6x^2 - 8x + 14}{(3x+4)(x-1)}$$

$$= \frac{-6x^2 + 7x - 1}{(3x+4)(x-1)}$$

$$= \frac{(-6x+1)(x-1)}{(3x+4)(x-1)}$$

$$f(x) = \frac{-6x+1}{3x+4}$$

$$b) \text{ Let } u = -6x + 1 \quad \text{Let } v = 3x + 4$$

$$\frac{du}{dx} = -6$$

$$\frac{dv}{dx} = 3$$



Question 2 continued

Quotient rule:

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$= \frac{(3x+4)(-6) - (-6x+1)(3)}{(3x+4)(3x+4)}$$

$$= \frac{-18x - 24 + 18x - 3}{(3x+4)^2}$$

$$= \frac{-27}{(3x+4)^2}$$



Leave
blank

Question 2 continued



Question 2 continued

(Total 7 marks)



3. (a) By writing $\operatorname{cosec} x$ as $\frac{1}{\sin x}$, show that

$$\frac{d(\operatorname{cosec} x)}{dx} = -\operatorname{cosec} x \cot x \quad (3)$$

Given that $y = e^{3x} \operatorname{cosec} 2x$, $0 < x < \frac{\pi}{2}$,

- (b) find an expression for $\frac{dy}{dx}$. (3)

The curve with equation $y = e^{3x} \operatorname{cosec} 2x$, $0 < x < \frac{\pi}{2}$, has a single turning point.

- (c) Show that the x -coordinate of this turning point is at $x = \frac{1}{2} \arctan k$ where the value of the constant k should be found. (2)

$$a) \frac{d}{dx} (\operatorname{cosec} x) = \frac{d}{dx} (\sin x)^{-1}$$

$$\text{Let } u = \sin x$$

$$\text{Let } y = u^{-1}$$

$$\frac{du}{dx} = \cos x$$

$$\frac{dy}{du} = -u^{-2}$$

Chain rule =

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$$

$$= -\frac{1}{\sin^2 x} \times \cos x$$

$$= -\frac{1}{\sin x} \times \frac{\cos x}{\sin x}$$

$$= -\operatorname{cosec} x \cot x$$



Question 3 continued

$$b) y = e^{3x} \operatorname{cosec} 2x$$

$$\text{Let } u = e^{3x}$$

$$\text{Let } v = \operatorname{cosec} 2x$$

$$\frac{du}{dx} = 3e^{3x}$$

$$\frac{dv}{dx} = -2 \operatorname{cosec} 2x \cot 2x$$

Product rule:

$$\frac{d}{dx}(uv) = v \frac{du}{dx} + u \frac{dv}{dx}$$

$$= \operatorname{cosec} 2x (3e^{3x}) + (e^{3x}) (-2 \operatorname{cosec} 2x \cot 2x)$$

$$= e^{3x} \operatorname{cosec} 2x (3 - 2 \cot 2x)$$

d) Turning point when

$$\frac{dy}{dx} = 0$$

$$(e^{3x}) (\operatorname{cosec} 2x) (3 - 2 \cot 2x) = 0$$

$$\text{Either } e^{3x} = 0$$

Reject as not valid.

$$\text{Or } \operatorname{cosec} 2x = 0$$

Reject as not valid

$$\text{Or } 3 - 2 \cot 2x = 0 \Rightarrow x = \frac{1}{2} \operatorname{arccot} \frac{3}{2}$$

$$\Rightarrow x = \frac{1}{2} \operatorname{arctan} \frac{2}{3}$$

(Total 8 marks)

Q3



4. A pot of coffee is delivered to a meeting room at 11am. At a time t minutes after 11am the temperature, $\theta^\circ\text{C}$, of the coffee in the pot is given by the equation

$$\theta = A + 60e^{-kt}$$

where A and k are positive constants.

Given also that the temperature of the coffee at 11am is 85°C and that 15 minutes later it is 58°C ,

Condition

- (a) find the value of A .


(1)

- (b) Show that $k = \frac{1}{15} \ln\left(\frac{20}{11}\right)$

(3)

- (c) Find, to the nearest minute, the time at which the temperature of the coffee reaches 50°C .

(4)



85°C is the initial temperature at 11am, when $t=0$.

$$\underline{A = 85 - 60 = 25^\circ\text{C}}$$

✓

b)

$$\theta = A + 60e^{-kt}$$

$$\theta - A = 60e^{-kt}$$

$$e^{-kt} = \frac{\theta - A}{60}$$

$$e^{-k(15)} = \frac{58 - 25}{60}$$



Question 4 continued

$$e^{-k(15)} = \frac{33}{60}$$

$$e^{+15k} = \frac{20}{11}$$

$$\ln e^{15k} = \ln\left(\frac{20}{11}\right)$$

Take logs.

$$15k(1) = \ln\left(\frac{20}{11}\right)$$

$$k = \frac{1}{15} \ln\left(\frac{20}{11}\right)$$

✓

$$c) \quad \theta = A + 60 e^{-kt}$$

Find t when $\theta = 50^\circ\text{C}$

$$50 = 25 + 60 e^{-kt}$$

$$e^{-kt} = \frac{25}{60}$$

$$e^{+kt} = \frac{12}{5}$$

$$\ln(e^{kt}) = \ln \frac{12}{5}$$

Taking logs.



Question 4 continued

$$kt = \ln \frac{12}{5}$$

$$t = \frac{1}{k} \ln \frac{12}{5}$$

$$= \frac{15}{\ln \left(\frac{20}{11} \right)} \ln \frac{12}{5}$$

↓

$$= 21.9659$$

$$\underline{t \approx 22.0 \text{ minutes}}$$

✓



Question 4 continued

Q4

13

P 4 3 1 3 6 A 0 1 3 2 8

5.




Figure 1

The curve shown in Figure 1 has equation

$$x = 3 \sin y + 3 \cos y, \quad -\frac{\pi}{4} < y < \frac{\pi}{4}$$

(a) Express the equation of the curve in the form

$$x = R \sin(y + \alpha), \text{ where } R \text{ and } \alpha \text{ are constants, } R > 0 \text{ and } 0 < \alpha < \frac{\pi}{2} \quad (3)$$

(b) Find the coordinates of the point on the curve where the value of $\frac{dy}{dx}$ is $\frac{1}{2}$.

Give your answers to 3 decimal places.

(6)

$$\begin{aligned} \text{a)} \quad 3 \sin y + 3 \cos y &= R \sin(y + \alpha) \\ &= R \sin y \cos \alpha + R \cos y \sin \alpha \end{aligned}$$

Comparing coefficients

$$R \cos \alpha = 3 \quad (1) \quad R \sin \alpha = 3 \quad (2)$$

$$\therefore (2) \div (1) \quad \tan \alpha = \frac{3}{3} = 1$$

$$\therefore \alpha = \frac{\pi}{4}$$



Question 5 continued

$$①^2 + ②^2 \quad R^2 = 3^2 + 3^2$$

$$\underline{R = \sqrt{18} = 3\sqrt{2}}$$

$$b) \quad x = R \sin(y + \alpha)$$

$$x = 3\sqrt{2} \sin(y + \pi/4)$$

$$\frac{dx}{dy} = 3\sqrt{2} \cos(y + \pi/4)$$

$$\text{If } \frac{dy}{dx} = \frac{1}{2}$$

$$\text{Then } \frac{dx}{dy} = 3\sqrt{2} \cos(y + \pi/4) = 2$$

$$\cos(y + \pi/4) = \frac{2}{3\sqrt{2}} = \frac{\sqrt{2}}{3}$$

$$y + \pi/4 = 1.0799$$

$$y = 0.2945 \quad \underline{\underline{\approx 0.295}} \quad (3 \text{ D.P.})$$

$$\Rightarrow x = 3\sqrt{2} \sin(0.2945 + \pi/4)$$

$$= 3\sqrt{2} (0.8819)$$

$$\underline{x = 3.742} \quad (3 \text{ P.P.})$$

$$\text{Coordinates: } (3.742, 0.295)$$



Leave
blank

Question 5 continued



Question 5 continued

(Total 9 marks)



6. Given that a and b are constants and that $0 < a < b$,

(a) on separate diagrams, sketch the graph with equation

(i) $y = |2x + a|$,

(ii) $y = |2x + a| - b$.

Show on each sketch the coordinates of each point at which the graph crosses or meets the axes.



(6)

(b) Solve, for x , the equation

$$|2x + a| - b = \frac{1}{3}x$$

giving any answers in terms of a and b .

(4)



Question 6 continued

$$\text{RHS: } y = 2x + a - b = \frac{1}{3}x$$

$$2x - \frac{1}{3}x = b - a$$

$$\frac{5}{3}x = b - a$$


$$x = \frac{3}{5}(b - a)$$

$$\text{LHS: } y = -2x - (a + b) = \frac{1}{3}x$$

$$-2\frac{1}{3}x = a + b$$

$$-\frac{7}{3}x = a + b$$

$$x = -\frac{3}{7}(a + b)$$



Leave
blank

Question 6 continued



Leave
blank**Question 6 continued****Q6****(Total 10 marks)**

7. (i) (a) Prove that

$$\cos 3\theta \equiv 4 \cos^3 \theta - 3 \cos \theta$$

(You may use the double angle formulae and the identity

$$\cos(A+B) \equiv \cos A \cos B - \sin A \sin B$$

(4)

(b) Hence solve the equation

$$2 \cos 3\theta + \cos 2\theta + 1 = 0$$

giving answers in the interval $0 \leq \theta \leq \pi$.

Solutions based entirely on graphical or numerical methods are not acceptable.

(6)

(ii) Given that $\theta = \arcsin x$ and that $0 < \theta < \frac{\pi}{2}$, show that

$$\cot \theta = \frac{\sqrt{1-x^2}}{x}, \quad 0 < x < 1$$

(3)

a) LHS:

$$\cos 3\theta = \cos(\theta + 2\theta)$$

$$= \cos \theta \cos 2\theta - \sin \theta \sin 2\theta$$

$$= \cos \theta (\cos^2 \theta - \sin^2 \theta) - \sin \theta 2 \sin \theta \cos \theta$$

$$= \cos^3 \theta + \cos \theta (- (1 - \cos^2 \theta)) - 2 \sin^2 \theta \cos \theta$$

$$= \cos^3 \theta - \cos \theta + \cos^3 \theta - 2(1 - \cos^2 \theta) \cos \theta$$

$$= 2 \cos^3 \theta - \cos \theta - 2 \cos \theta + 2 \cos^3 \theta$$

$$= \underline{4 \cos^3 \theta - 3 \cos \theta}$$

Using:

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$



Question 7 continued

$$b) 2 \cos 3\theta + \cos 2\theta + 1 = 0$$

$$2(4\cos^3\theta - 3\cos\theta) + (\cos^2\theta - \sin^2\theta) + 1 = 0$$

$$8\cos^3\theta - 6\cos\theta + \cos^2\theta - (1 - \cos^2\theta) + 1 = 0$$

$$8\cos^3\theta + 2\cos^2\theta - 6\cos\theta = 0$$

$$2\cos\theta (4\cos^2\theta + \cos\theta - 3) = 0$$

$$(2\cos\theta)(\cos\theta + 1)(4\cos\theta - 3) = 0$$

$$\text{Either } 2\cos\theta = 0$$

$$\text{Or } \cos\theta = -1$$

$$\cos\theta = 0$$

$$\theta = \pi \quad \checkmark$$

$$\theta = \frac{\pi}{2} \quad \checkmark$$

$$\text{Or } 4\cos\theta = 3$$

$$\cos\theta = \frac{3}{4}$$

$$\theta = 0.723 \text{ rad.} \quad \checkmark$$

$$\text{Range } 0 \leq \theta \leq \pi$$

$$d) \theta = \arcsin x$$

$$\Rightarrow x = \sin\theta$$

$$\cot\theta = \frac{1}{\tan\theta} = \frac{\cos\theta}{\sin\theta}$$

$$= \frac{\sqrt{\cos^2\theta}}{\sin\theta} = \frac{\sqrt{1 - \sin^2\theta}}{\sin\theta}$$

$$\therefore \cot\theta = \frac{\sqrt{1 - x^2}}{x}, \quad 0 < x < 1$$



Question 7 continued

Q7

8. The function f is defined by

$$f: x \rightarrow 3 - 2e^{-x}, \quad x \in \mathbb{R}$$

(a) Find the inverse function, $f^{-1}(x)$ and give its domain.

(5)

(b) Solve the equation $f^{-1}(x) = \ln x$.

(4)

The equation $f(t) = ke^t$, where k is a positive constant, has exactly one real solution.

(c) Find the value of k .

(4)

$$f: x \rightarrow 3 - 2e^{-x} \quad \text{Range } f(x) < 3$$

$$y = 3 - 2e^{-x}$$

$$2e^{-x} = 3 - y$$

$$e^{-x} = \frac{3-y}{2}$$

$$\ln e^{-x} = \ln \left(\frac{3-y}{2} \right)$$

$$x = -\ln \left(\frac{3-y}{2} \right)$$

$$x = \ln \left(\frac{2}{3-y} \right) \quad \text{Dom}$$

$$\therefore f^{-1}(x) = \ln \left(\frac{2}{3-x} \right) \quad \checkmark \quad \text{Domain of inverse} = \text{range of function.}$$

$$\text{So } x < 3$$



Question 8 continued

$$b) f^{-1}(x) = \ln x$$

$$\therefore \ln\left(\frac{2}{3-x}\right) = \ln x$$

$$\frac{2}{3-x} = x$$

$$2 = x(3-x)$$

$$2 = 3x - x^2$$

$$x^2 - 3x + 2 = 0$$


$$(x-2)(x-1) = 0$$

$$\text{Either } x = 2$$

$$\text{Or } x = 1$$



$$c) f(t) = k e^t$$



$$\therefore 3 - 2e^{-t} = k e^t$$

$$k e^t + 2e^{-t} = 3$$



Question 8 continued

$$ke^t + 2e^t - 3 = 0$$

$$ke^{2t} - 3e^t + 2 = 0 \quad (xe^t)$$

$$ax^2 + bx + c = 0$$

Using

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$1 \text{ real solution so } b^2 - 4ac = 0$$

$$(-3)^2 = 4k2$$

$$k = \frac{9}{8} \quad \checkmark$$

Q8

(Total 13 marks)

TOTAL FOR PAPER: 75 MARKS

END

