

Write your name here	
Surname MODEL ANSWERS	Other names
Pearson Edexcel International Advanced Level	Centre Number Candidate Number
Further Pure Mathematics FP1 Advanced/Advanced Subsidiary	
Wednesday 29 January 2014 – Morning Time: 1 hour 30 minutes	Paper Reference 6667A/01
You must have: Mathematical Formulae and Statistical Tables (Pink)	Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P43019A

©2014 Pearson Education Ltd.
5/5/5/5/



PEARSON

1. $f(x) = 2x - 5 \cos x$, where x is in radians.

(a) Show that the equation $f(x) = 0$ has a root α in the interval $[1, 1.4]$.

(2)

(b) Starting with the interval $[1, 1.4]$, use interval bisection twice to find an interval of width 0.1 which contains α .

(3)

a) $f(1) = 2(1) - 5 \cos(1) = -0.702$

$$f(1.4) = 2(1.4) - 5 \cos(1.4) = +1.950$$

Due to the change in sign, where $f(x)$ is a continuous function, there must be a root in the interval $[1, 1.4]$.

b) Midpoint of interval $[1, 1.4] = 1.2$

$$f(1.2) = 0.588$$

By change of sign rule, new interval: $[1, 1.2]$

Midpoint of interval $[1, 1.2] = 1.1$

$$f(1.1) = -0.0680$$

By change of sign rule, new interval: $[1.1, 1.2]$



Leave
blank

Question 1 continued

This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There is no handwriting or other markings on the paper.

Q1

(Total 5 marks)



2.

(i) $A = \begin{pmatrix} -4 & 10 \\ -3 & k \end{pmatrix}$, where k is a constant.

The triangle T is transformed to the triangle T' by the transformation represented by A .

Given that the area of triangle T' is twice the area of triangle T ,
find the possible values of k .

(4)

(ii) Given that

$$B = \begin{pmatrix} 1 & -2 & 3 \\ -2 & 5 & 1 \end{pmatrix}, \quad C = \begin{pmatrix} 2 & 8 \\ 0 & 2 \\ 1 & -2 \end{pmatrix}$$

find BC .

(3)

(i) Area of image = Area of object $\times |\det A|$

As Area $T' = 2$ Area T

$|\det A| = 2$ (modulus)

$\therefore -4k - (-3 \times 10) = 2$ Or

$-4k + 30 = 2$ $-(-4k + 30) = 2$

$4k = -28$

$k = -7$

$4k = 32$

$k = 8$



Question 2 continued

B C

$$= \begin{bmatrix} 1 & -2 & 3 \\ -2 & 5 & 1 \end{bmatrix} \begin{bmatrix} 2 & 8 \\ 0 & 2 \\ 1 & -2 \end{bmatrix} = \begin{bmatrix} 2+3 & 8-4-6 \\ -4+1 & -16+10-2 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & -2 \\ -3 & -8 \end{bmatrix}$$

Q2

(Total 7 marks)




3. A rectangular hyperbola has parametric equations

$$x = 2t, \quad y = \frac{2}{t}, \quad t \neq 0$$

Points P and Q on this hyperbola have parameters $t = \frac{1}{2}$ and $t = 4$ respectively.

The line L , which passes through the origin O , is perpendicular to the chord PQ .

Find an equation for L .



$$\begin{aligned} \text{Gradient } m_{PQ} &= \frac{y_2 - y_1}{x_2 - x_1} = \frac{\frac{1}{2} - 4}{8 - 1} \\ &= -\frac{7}{2} \times \frac{1}{7} = \underline{\underline{-\frac{1}{2}}} \end{aligned}$$

$$\therefore \text{Line } L: y = mx + c$$

$$y = -\frac{1}{2}x + c$$

$$\text{Using } (0, 0) \quad 0 = -\frac{1}{2}(0) + c \Rightarrow c = 0$$

$$\therefore \text{Line } L: \underline{\underline{y = -\frac{1}{2}x}}$$



Question 3 continued

Leave
blank

Q3

(Total 4 marks)



4. $f(x) = 2x^{\frac{1}{2}} - \frac{6}{x^2} - 3, \quad x > 0$

A root β of the equation $f(x) = 0$ lies in the interval $[3, 4]$.

Taking 3.5 as a first approximation to β , apply the Newton-Raphson process once to $f(x)$ to obtain a second approximation to β . Give your answer to 3 decimal places.

(5)

$$f(x) = 2x^{\frac{1}{2}} - 6x^{-2} - 3$$

$$f'(x) = \left(\frac{1}{2}\right)2x^{-\frac{1}{2}} - (-2)6x^{-3} + 0$$

$$f'(x) = x^{-\frac{1}{2}} + 12x^{-3}$$

Newton-Raphson:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$= 3.5 - \frac{(2(3.5)^{\frac{1}{2}} - 6(3.5)^{-2} - 3)}{(3.5)^{-\frac{1}{2}} + 12(3.5)^{-3}}$$

$$x_{n+1} = 3.190742075 \approx 3.191 \quad (3dp)$$

$$x_{n+2} = 3.208835389 \approx \underline{\underline{3.209}} \quad (3dp)$$



Question 4 continuedLeave
blank**(Total 5 marks)****Q4**

5.

$$z = 5 + i\sqrt{3}, \quad w = \sqrt{3} - i$$

(a) Find the value of $|w|$.

(1)

Find in the form $a + ib$, where a and b are real constants,(b) zw , showing clearly how you obtained your answer,

(2)

(c) $\frac{z}{w}$, showing clearly how you obtained your answer.

(3)

Given that

$$\arg(z + \lambda) = \frac{\pi}{3}, \quad \text{where } \lambda \text{ is a real constant,}$$

(d) find the value of λ .

(2)

$$a) \quad |w| = \sqrt{(\sqrt{3})^2 + (-1)^2} = \sqrt{4} = \underline{\underline{2}}$$

✓

$$\begin{aligned} b) \quad zw &= (5 + i\sqrt{3})(\sqrt{3} - i) \\ &= 5\sqrt{3} - 5i + 3i - i^2\sqrt{3} \\ &= 5\sqrt{3} - 2i - (-1)\sqrt{3} \\ &= \underline{\underline{6\sqrt{3} - 2i}} \end{aligned}$$

✓


$$\begin{aligned} c) \quad \frac{z}{w} &= \frac{(5 + i\sqrt{3})}{(\sqrt{3} - i)} \times \frac{(\sqrt{3} + i)}{(\sqrt{3} + i)} \\ &= \frac{5\sqrt{3} + 5i + 3i + \sqrt{3}i^2}{3 + 3i - 3i - i^2} \\ &= \frac{5\sqrt{3} + 8i - \sqrt{3}}{3 + 1} = \frac{5\sqrt{3} - \sqrt{3} + 8i}{4} \\ &= \underline{\underline{\sqrt{3} + 2i}} \end{aligned}$$

✓



Question 5 continued

$$d) \arg(z + 1) = \frac{\pi}{3}$$



$$\text{If } z = x + iy, \tan \theta = \frac{y}{x}$$

$$\text{If } \tan \theta = \frac{\pi}{3}$$

$$\theta = \sqrt{3}$$

$$\text{If, } z = 5 + i\sqrt{3}$$

$$\therefore z + 1 = 5 + 1 + i\sqrt{3}$$

$$\therefore \frac{y}{x} = \frac{\sqrt{3}}{5+1} = \sqrt{3} \Rightarrow \underline{\underline{\lambda = -4}}$$



Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5.



Question 5 continuedLeave
blank

Q5

(Total 8 marks)

P 4 3 0 1 9 A 0 1 3 3 2

6. (a) Use the standard results for $\sum_{r=1}^n r^3$ and $\sum_{r=1}^n r$ to show that for all positive integers n ,

$$\sum_{r=1}^n r(r+1)(r-1) = \frac{1}{4}n(n+1)(n-1)(n+a)$$

where a is an integer to be determined.

(4)

- (b) Hence find the value of n , where $n > 1$, that satisfies

$$\sum_{r=1}^n r(r+1)(r-1) = 10 \sum_{r=1}^n r^2$$

(5)

$$a) \sum_{r=1}^n r^3 = \frac{n^2}{4} (n+1)^2$$

$$\sum_{r=1}^n r = \frac{n}{2} (n+1)$$

$$\sum_{r=1}^n r(r+1)(r-1) = \sum_{r=1}^n r(r^2 - 1)$$

$$= \sum_{r=1}^n r^3 - \sum_{r=1}^n r$$

$$= \frac{n^2}{4} (n+1)^2 - \frac{n}{2} (n+1)$$

$$= \frac{n^4}{4} + \frac{2n^3}{4} + \frac{n^2}{4} - \frac{n^2}{2} - \frac{n}{2}$$

$$= \frac{n^4}{4} + \frac{2n^3}{4} - \frac{n^2}{4} - \frac{2n}{4}$$

$$= \frac{1}{4} n(n+1)(n-1)(n+2)$$



$$n(n^2-1)(n-2) = (n^3-n)(n-2) = n^4 - 2n^3 - n^2 + 2n$$

Question 6 continued

or/

$$= \frac{1}{4}n(n+1)[n(n+1)-2] \text{ factorise}$$

$$= \frac{1}{4}n(n+1)[n^2+n-2] \text{ simplify}$$

$$= \frac{1}{4}n(n+1)(n-1)(n+2)$$

$$= \frac{1}{4}n(n+1)(n-1)(n+2)$$

$$\Rightarrow \underline{a = 2}$$



This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is no text or other markings on the paper.

7. $\mathbf{P} = \begin{pmatrix} 3a & -2a \\ -b & 2b \end{pmatrix}, \quad \mathbf{M} = \begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix}$

where a and b are non-zero constants.

(a) Find $\mathbf{P}^{-1}$, leaving your answer in terms of a and b .

(3)

Given that

$$\mathbf{M} = \mathbf{PQ}$$

(b) find the matrix $\mathbf{Q}$, giving your answer in its simplest form.

(3)

$$\begin{aligned} \text{a) } \det \underline{\mathbf{P}} &= 3a \times 2b - (-b) \times (-2a) \\ &= 6ab - 2ab = 4ab \end{aligned}$$

$$\begin{aligned} \text{cofactors } \underline{\mathbf{C}} &= \begin{pmatrix} 2b & +b \\ +2a & +3a \end{pmatrix} & \underline{\mathbf{C}}^T &= \begin{pmatrix} 2b & 2a \\ +b & +3a \end{pmatrix} \text{ transpose} \\ & \begin{pmatrix} + & - \\ - & + \end{pmatrix} \end{aligned}$$

$$\mathbf{P}^{-1} = \frac{1}{4ab} \begin{pmatrix} 2b & 2a \\ b & 3a \end{pmatrix}$$

$$\text{b) } \underline{\mathbf{M}} = \underline{\mathbf{P}} \underline{\mathbf{Q}}$$

$$\underline{\mathbf{P}}^{-1} \underline{\mathbf{M}} = \underline{\mathbf{P}}^{-1} \underline{\mathbf{P}} \underline{\mathbf{Q}}$$

$$\underline{\mathbf{P}}^{-1} \underline{\mathbf{M}} = \underline{\mathbf{I}} \underline{\mathbf{Q}}$$

$$\underline{\mathbf{Q}} = \underline{\mathbf{P}}^{-1} \underline{\mathbf{M}}$$



Question 7 continued

$$\underline{Q} = \frac{1}{4ab} \begin{pmatrix} 2b & 2a \\ b & -3a \end{pmatrix} \begin{pmatrix} -6a & 7a \\ 2b & -b \end{pmatrix}$$

$$= \frac{1}{4ab} \begin{pmatrix} -12ab + 4ab & 14ab - 2ab \\ -6ab + 6ab & 7ab - 3ab \end{pmatrix}$$

$$= \frac{1}{4ab} \begin{pmatrix} -8ab & 12ab \\ 0 & 4ab \end{pmatrix}$$

$$= \begin{pmatrix} -2 & 3 \\ 0 & 1 \end{pmatrix}$$

Q7

(Total 6 marks)



P 4 3 0 1 9 A 0 1 9 3 2

8. The parabola C has equation $y^2 = 4ax$, where a is a positive constant.

The point $P(ap^2, 2ap)$ lies on the parabola C .

- (a) Show that an equation of the normal to C at P is


$$y + px = ap^3 + 2ap \quad (5)$$

The normal to C at the point P meets the x -axis at the point $(6a, 0)$ and meets the directrix of C at the point D . Given that $p > 0$,

- (b) find, in terms of a , the coordinates of D . (4)

Given also that the directrix of C cuts the x -axis at the point X ,

- (c) find, in terms of a , the area of the triangle XPD , giving your answer in its simplest form. (3)



$$x = ap^2 \quad y = 2ap$$

$$\frac{dx}{dp} = 2ap$$

$$\frac{dy}{dp} = 2a$$

$$\therefore \frac{dy}{dx} = \frac{dy}{dp} \times \frac{dp}{dx}$$

$$\frac{dy}{dx} = \frac{2a}{2ap} = \frac{1}{p} = m_T$$



Question 8 continued

$$m_N = -\frac{1}{m_T}$$

$$m_N = -p$$

$$y - y_1 = m(x - x_1)$$

$$\text{At } P(ap^2, 2ap)$$

Normal:

$$y - 2ap = -p(x - ap^2)$$

$$y + px = ap^3 + 2ap$$

$$\text{b) At } (6a, 0)$$

$$0 + p(6a) = ap^3 + 2ap$$

$$6ap - 2ap = ap^3$$

$$4p = p^3$$

$$p^2 = 4$$

$$\therefore p = 2$$

$$\therefore y + 2x = 8a + 4a = 12a \quad \checkmark$$


Normal intersects directrix at D when $x = -a$

$$y + 2(-a) = 12a$$

$$y = 14a \Rightarrow D(-a, 14a) \quad \checkmark$$



Question 8 continued



$$\begin{aligned}
 \text{Area } \triangle xPD &= \frac{1}{2} b \times h \\
 &= \frac{1}{2} \times 14a \times (4a + a) \\
 &= \underline{35a^2} \quad \text{units}^2
 \end{aligned}$$



Question 8 continuedLeave
blank**(Total 12 marks)****Q8**

9. Given that $z = x + iy$, where $x \in \mathbb{R}, y \in \mathbb{R}$, find the value of x and the value of y such that

$$(3 - i)z^* + 2iz = 9 - i$$

where z^* is the complex conjugate of z .

(8)

$$z^* = x - iy$$

$$(3 - i)z^* = (3 - i)(x - iy)$$

$$= 3x - 3iy - ix + i^2y$$

$$= 3x - y - (3y + x)i$$

$$\therefore (3 - i)z^* + 2iz$$

$$= 3x - y - (3y + x)i + 2i(x + iy)$$

$$= 3x - y - (3y + x)i + 2ix + 2i^2y$$

$$= 3x - 3y + (x - 3y)i = 9 - i$$

Equating real + imaginary parts:

$$\text{Real: } 3x - 3y = 9 \quad \text{Imaginary: } x - 3y = -1 \quad \textcircled{2}$$

$$\textcircled{1} - \textcircled{2}$$

$$2x + 0 = 10$$

$$\underline{x = 5}$$

Sub into $\textcircled{1}$

$$3y = 3(5) - 9$$

$$3y = 6$$

$$\underline{y = 2}$$



Question 9 continuedLeave
blank

P 4 3 0 1 9 A 0 2 5 3 2

Leave
blank

Question 9 continued

Lined area for writing the answer to Question 9.



Question 9 continuedLeave
blank**(Total 8 marks)****Q9**

10. (i) A sequence of numbers $u_1, u_2, u_3, \dots$, is defined by

$$u_{n+1} = 5u_n + 3, \quad u_1 = 3$$

Prove by induction that, for $n \in \mathbb{Z}^+$,

$$u_n = \frac{3}{4}(5^n - 1) \quad (5)$$

- (ii) Prove by induction that, for $n \in \mathbb{Z}^+$,

$$f(n) = 5(5^n) - 4n - 5 \text{ is divisible by } 16. \quad (6)$$

(i)

$$\text{Let } n=1, \quad u_1 = \frac{3}{4}(5^1 - 1)$$

$$u_1 = \frac{3}{4}(4) = 3$$

Therefore this is true when $n=1$.

Assumption - True for $n=k$

$$\therefore u_k = \frac{3}{4}(5^k - 1)$$

Prove for $n=k+1$

$$u_{k+1} = 5u_k + 3$$

$$= 5 \left[\frac{3}{4}(5^k - 1) \right] + 3$$

$$= \frac{3}{4}(5^1 \times 5^k) - \frac{15}{4} + 3$$

$$= \frac{3}{4}5^{k+1} - \frac{3}{4}$$

$$u_{k+1} = \frac{3}{4}(5^{k+1} - 1)$$



Question 10 continued

The statement is true for $n = k+1$,
 true for $n = 1$
 on basis of assumption for $n = k$

$\Rightarrow$ True for $n \in \mathbb{Z}^+$

(ii) Prove, $f(n) = 5(5^n) - 4n - 5$ is divisible
 by 16

For $n=1$, $f(1) = 5(5^1) - 4(1) - 5 = 16$

therefore $f(1)$ is divisible by 16, this is true.

Assumption - true for $n=k$

$$f(k) = 5(5)^k - 4k - 5 \text{ is divisible by 16}$$

Prove true for $n = k+1$

$$\begin{aligned} f(k+1) &= 5(5)^{k+1} - 4(k+1) - 5 \\ &= 5 \times 5^1 \times 5^k - 4k - 4 - 5 \\ &= 5 \times 5(5)^k - (20k + 16k) + (-5 - 4 + 20) \\ &= 5[5(5)^k - 4k - 5] + 16k + 16 \end{aligned}$$

$$f(k+1) = 5f(k) + 16k + 16$$

If $f(k)$ is divisible by 16, then so is $5f(k)$
 then so is $5f(k) + 16k + 16$

True for $n=k+1$ so true for $n \in \mathbb{Z}^+$



This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is no text or other markings on the paper.

Question 10 continuedLeave
blank

P 4 3 0 1 9 A 0 3 1 3 2

Q10

[illegible]**TOTAL FOR PAPER: 75 MARKS**

32

